

UNCLASSIFIED

AD

426482

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

nato

ORC 63-27 (RR)
21 OCTOBER 1963

426482

426482

BOUNDS ON INTEGRALS WITH
APPLICATIONS TO RELIABILITY
PROBLEMS

by

Richard E. Barlow


CATALOGED BY DDC

AS AD No. _____

OPERATIONS RESEARCH CENTER

INSTITUTE OF ENGINEERING RESEARCH

UNIVERSITY OF CALIFORNIA - BERKELEY



BOUNDS ON INTEGRALS WITH APPLICATIONS TO RELIABILITY PROBLEMS

by

Richard E. Barlow
Operations Research Center
University of California, Berkeley

21 October 1963

ORC 63-27 (RR)

This research has been partially supported by the Office of Naval Research under Contract Nonr-222(83) with the University of California. Reproduction in whole or in part is permitted for any purpose of the United States Government.

BOUNDS ON INTEGRALS WITH APPLICATIONS TO RELIABILITY PROBLEMS

1. Summary

Certain statistical problems in life testing and reliability lead to the consideration of minimizing and maximizing certain integrals of the form

$$\int_0^{\infty} \phi(x, \bar{F}(x)) dx \quad \bar{F}(x) = 1 - F(x)$$

under the assumption that the distribution function F has increasing hazard rate and that one or two moments are given. The minimizing (maximizing) distributions for the special cases considered are members of the extremal class considered by A. Marshall and the author [1]. Bounds on the expected values of the minimum and maximum order statistics as well as the operating characteristics of replacement policies based on age are obtained under these assumptions.

2. Statistical Motivation

For many statistical problems in life testing and reliability it is natural to assume that the life distribution F ($F(0^-) = 0$) has increasing hazard rate (is IHR). If F has density f , then F is IHR if and only if $f(x)/[1 - F(x)]$ is nondecreasing in x . In general F is said to be IHR if and only if $\ln[1 - F(x)]$ is concave where finite. This property is enjoyed by most of the commonly used life distributions; e.g.,

$$\text{Weibull: } f(x) = \begin{cases} \lambda \alpha x^{\alpha-1} \exp(-\lambda x^{\alpha}) & x \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$

for $\alpha \geq 1$, $\lambda > 0$, and

$$\text{Gamma: } f(x) = \begin{cases} \frac{\lambda^\alpha x^{\alpha-1} \exp(-\lambda x)}{\Gamma(\alpha)} & x \geq 0 \\ 0 & \text{elsewhere} \end{cases}$$

for $\alpha \geq 1$, $\lambda > 0$. Distributions F for which $\ln[1 - F(x)]$ is convex on $[0, \infty)$ are called DHR for decreasing hazard rate. Properties of distributions with monotone hazard rate are discussed in [2].

In addition to this assumption, some prior information in the form of the mean μ_1 , and the variance $\sigma^2 = \mu_2 - \mu_1^2$ of the distribution may be known. Since in the IHR case

$$\frac{\sigma}{\mu_1} \leq 1$$

with equality only for exponential distribution, an estimate of the coefficient of variation σ/μ_1 may represent the statistician's belief in the amount of departure of the failure distribution from exponentiality.

With the mean and variance assumed known we consider the problem of minimizing and maximizing certain integrals of the form

$$(2.1) \quad \int_0^\infty \phi(x, \bar{F}(x)) dx$$

where F is IHR. Integrals of such functions have been considered by Karlin [3], Rustagi [4] and others for general distributions.

(See [4] for additional references). A related but more specialized problem is that of minimizing (maximizing) integrals of the form

$$\int_0^\infty \rho(x) dF(x)$$

assuming that certain moments of F are specified, (cf. (3.1) and Theorem 4.1). Assuming no conditions other than the moment conditions, this problem has been solved by H. P. Mulholland and C. A. Rogers (1958). Their methods as well as those of many others (e.g. Karlin and Shapley (1953)) use the fact that the class of all distributions subject to specified moment conditions are closed under convex combinations. However, the class of IHR distributions is not closed under convex combinations [2, p. 381] and, therefore, we cannot use the classical methods. Instead we consider very special cases using the methods of [1]. However, we do obtain fairly explicit bounds which can be easily machine computed.

Integrals such as (2.1) often arise in statistical applications. For example, let $U_1 \leq U_2 \leq \dots \leq U_n$ denote n ordered observations from a population with IHR cdf F . Let $W_n = U_n - U_1$ denote the sample range. Then we may seek to minimize (maximize)

$$E(W_n) = \int_0^{\infty} [1 - F(x)^n - (\bar{F}(x))^n] dx$$

where $\bar{F}(x) = 1 - F(x)$. The integrand is of the form $\phi(x, y)$ where

$$\phi(x, y) = 1 - (1 - y)^n - y^n$$

is concave in y .

For another example, consider a replacement policy of the form: Replace at time t or at failure, whichever occurs first. We may then seek to minimize (maximize) the following integrals:

$$(i) \int_0^t \bar{F}(x) dx \quad \bar{F}(x) = 1 - F(x)$$

i.e., the expected time between removals for either a failed or nonfailed item.

$$(ii) \quad \frac{\int_0^t \bar{F}(x) dx}{F(t)}$$

i.e., the expected time between failure removals using a replacement policy based on t . The reciprocal of this quantity occurs as an upper bound on the renewal function divided by t , [5].

$$(iii) \quad \frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)}$$

i.e., the expected time between planned replacements using a replacement policy based on t .

A related expression is:

$$(iv) \quad \int_t^\infty \bar{F}(x) dx / \bar{F}(t)$$

i.e., the mean residual life of an item aged t . Bounds on densities and failure rate functions will be discussed in a future joint paper with A. Marshall.

3. Bounds on Integrals when F is IHR with Specified Mean

If $\phi(x, y)$ is sufficiently smooth the problem of minimizing (maximizing)

$$\int_0^\infty \phi(x, \bar{F}(x)) dx$$

is comparatively simple. If F is IHR with mean μ_1 and $\rho(x)$

is increasing (decreasing) then

$$(3.1) \quad \int_0^{\mu_1} \rho(x) dx \underset{(\sum)}{\leq} \int_0^{\infty} \rho(x) \bar{F}(x) dx \underset{(\sum)}{\leq} \int_0^{\infty} \rho(x) e^{\frac{x}{-\mu_1}} dx$$

This result is essentially due to Karlin and Novikoff and follows easily from their method of proof in [6]. Using (3.1) and the method of proof in Karlin [3] and Rustagi [4] we obtain

THEOREM 3.1: If

- (i) F is IHR, $F(0) = 0$,
- (ii) $\int_0^{\infty} x dF(x) = \mu_1$,
- (iii) $\phi(x, y)$ is convex (concave) in y
- (iv) $\frac{\partial}{\partial y} \phi(x, y)$ is nonincreasing (nondecreasing) in x ,

then

$$\int_0^{\infty} \phi(x, \bar{G}_{\mu_1}(x)) dx \underset{(\sum)}{\geq} \int_0^{\infty} \phi(x, \bar{F}(x)) dx \underset{(\sum)}{\geq} \int_0^{\infty} \phi(x, e^{\frac{x}{-\mu_1}}) dx$$

when the indicated integrals exist and where G_{μ_1} is the distribution degenerate at μ_1 .

PROOF: We show the lower bound, assuming $\phi(x, y)$ is convex in y and $\frac{\partial}{\partial y} \phi(x, y)$ nonincreasing in x . Let $\bar{G}_0(x) = e^{-x/\mu_1}$ and define

$$(3.2) \quad I(\lambda) = \int_0^{\infty} \phi[x, \lambda \bar{G}_0(x) + (1-\lambda) \bar{F}(x)] dx \quad \text{for } 0 \leq \lambda \leq 1.$$

Using (iii) it is easy to check that $I(\lambda)$ is convex in λ . If $I(\lambda)$ achieves its minimum at $\lambda = 1$ then certainly $\bar{G}_0(x)$ will minimize (2.1). This is possible if and only if

$$I'(\lambda) \big|_{\lambda=1} \leq 0.$$

Now

$$I'(\lambda) \big|_{\lambda=1} = \int_0^\infty \frac{\partial}{\partial y} \phi(x, \bar{G}_0(x)) [\bar{G}_0(x) - \bar{F}(x)] dx.$$

By (iv), $\frac{\partial}{\partial y} \phi(x, y)$ is nonincreasing in x . By assumption (iii) $\frac{\partial}{\partial y} \phi(x, y)$ is nondecreasing in y . But if we let

$$y = \bar{G}_0(x), \text{ then}$$

$$\frac{\partial}{\partial y} \phi(x, y) \big|_{y=\bar{G}_0(x)} \text{ is nonincreasing in } x.$$

Therefore, by (3.1)

$$\begin{aligned} & \int_0^\infty \frac{\partial}{\partial y} \phi(x, y) \big|_{y=\bar{G}_0(x)} \bar{F}(x) dx \\ & \geq \int_0^\infty \frac{\partial}{\partial y} \phi(x, y) \big|_{y=\bar{G}_0(x)} \bar{G}_0(x) dx \end{aligned}$$

and hence $I'(\lambda) \big|_{\lambda=1} \leq 0$. Therefore

$$\int_0^\infty \phi(x, \bar{F}(x)) dx \geq \int_0^\infty \phi(x, \bar{G}_0(x)) dx.$$

If we assume $\phi(x, y)$ is concave in y and $\frac{\partial}{\partial y} \phi(x, y)$ is nondecreasing in x , then all inequalities are reversed.

The other inequalities follow in a similar manner. ||

Of course an analogous theorem can be stated for DHR distributions.

Let X_1, X_2, \dots, X_n denote n independent observations on a random variable X with IHR distribution F . Let $U_1 \leq U_2 \leq \dots \leq U_n$ denote the associated order statistics. Then

$$\begin{aligned} E[U_n - U_1] &= \int_0^\infty \{1 - [\bar{F}(x)]^n - [1 - \bar{F}(x)]^n\} dx \\ &= \int_0^\infty \phi(x, \bar{F}(x)) dx \end{aligned}$$

where $\phi(x, y) = 1 - y^n - (1 - y)^n$. Since ϕ is concave in y , Theorem 3.1 applies and we have

$$0 \leq E[U_n - U_1] \leq \mu_1 \sum_{k=1}^{n-1} 1/k$$

for sharp upper and lower bounds on the expected range. Intuitively, this inequality tells us that observations on IHR random variables tend to cluster together more closely than observations from an exponentially distributed random variable with the same mean. In a similar manner we obtain sharp upper and lower bounds on the expected values of the minimum and the maximum observations.

$$(3.2) \quad \frac{\mu_1}{n} \leq E[U_1] = \int_0^\infty [\bar{F}(x)]^n dx \leq \mu_1$$

$$(3.3) \quad \mu_1 \leq E[U_n] = \int_0^\infty [1 - F^n(x)] dx \leq \mu_1 \sum_{k=1}^n 1/k$$

Note that $\phi(x, y) = \alpha x^{\alpha-1} [1 - (1 - y)^n]$ satisfies the hypotheses of Theorem 3.1 for $\phi(x, y)$ concave in y and $\frac{\partial}{\partial y} \phi(x, y)$ non-decreasing in x when $\alpha \geq 1$. Using this ϕ we can bound all moments of U_n . Namely

$$\begin{aligned}
& \int_0^{\infty} \alpha x^{\alpha-1} [1 - G_{\mu_1}^n(x)] dx \leq E[U_n^\alpha] \\
& = \int_0^{\infty} \alpha x^{\alpha-1} [1 - F^n(x)] dx \leq \int_0^{\infty} \alpha x^{\alpha-1} [1 - G_0^n(x)] dx.
\end{aligned}$$

Inequalities (3.2) and (3.3) can be extended to include non-identically distributed random variables. The following proof is due to A. Marshall.

THEOREM 3.2: If $F_i(x)$ is IHR (DHR) with mean μ_i and $\bar{G}_i(x) = e^{-x/\mu_i}$ ($i = 1, 2, \dots, n$), then

$$(i) \quad \int_0^{\infty} \prod_{i=1}^n \bar{F}_i(x) dx \underset{(\leq)}{\geq} 1 / \sum_{i=1}^n 1/\mu_i$$

$$(ii) \quad \int_0^{\infty} [1 - \prod_{i=1}^n F_i(x)] dx \underset{(\sum)}{\leq} \int_0^{\infty} [1 - \prod_{i=1}^n G_i(x)] dx$$

PROOF: First we show (i). By (3.1)

$$\begin{aligned}
& \int_0^{\infty} \left[\prod_{j=1}^{i-1} \bar{F}_j(x) \prod_{j=i+1}^n \bar{G}_j(x) \right] \bar{F}_i(x) dx \\
& \underset{(\leq)}{\geq} \int_0^{\infty} \left[\prod_{j=1}^{i-1} \bar{F}_j(x) \prod_{j=i+1}^n \bar{G}_j(x) \right] \bar{G}_i(x) dx
\end{aligned}$$

for $1 \leq i \leq n$. By recursion we obtain

$$\int_0^{\infty} \prod_{j=1}^n \bar{F}_j(x) dx \underset{(\leq)}{\geq} \int_0^{\infty} \prod_{j=1}^n \bar{G}_j(x) dx = 1 / \sum_{i=1}^n \frac{1}{\mu_i}$$

To show (ii), note that again by (3.1)

$$\int_0^{\infty} \prod_{j=1}^{i-1} F_j(x) \prod_{j=i+1}^n G_j(x) \bar{F}_i(x) dx$$

$$\leq \int_0^{\infty} \prod_{j=1}^{i-1} F_j(x) \prod_{j=i+1}^n G_j(x) \bar{G}_i(x) dx$$

which in turn implies

$$\int_0^{\infty} [1 - \prod_{j=1}^i F_j(x) \prod_{j=i+1}^n G_j(x)] dx$$

$$\leq \int_0^{\infty} [1 - \prod_{j=1}^i F_j(x) \prod_{j=i}^n G_j(x)] dx$$

(ii) follows by recursion. ||

Part (i) tells us that the mean life of a series system with IHR components whose means are μ_i ($i=1, 2, \dots, n$) exceeds the mean life of a series system with exponentially distributed components and means μ_i ($i=1, 2, \dots, n$). However, just the reverse is true for parallel systems.

Additional properties of order statistics from IHR (DHR) distributions together with applications to life testing will appear in a future report.

4. Bounds on Integrals when F is IHR with Specified Mean and Variance

Assume now that F is IHR with mean μ_1 and μ_2 specified. In [1] the class of extremal distributions for bounding F were determined. Two rather special members of this class are

$$1 - G_{T_1}(x) = \begin{cases} \exp(-a_0 x) & x < T_1 \\ 0 & x \geq T_1 \end{cases}$$

and

$$1 - G_{T_0}(x) = \begin{cases} 1 & x \leq 1 - \sqrt{\mu_2 - 1} = T_0 \\ \exp\left[\frac{-(x - T_0)}{\sqrt{\mu_2 - 1}}\right] & x \geq T_0 \end{cases},$$

where a_0 and T_1 are chosen to satisfy the moment conditions. With these distributions and different assumptions on $\rho(x)$ we can improve the bounds in (3.1). The method we now use has been employed by Karlin and Novikoff [6] among others.

THEOREM 4.1 If

- (i) F is IHR, $F(0) = 0$
- (ii) $\mu_1 = 1$ and μ_2 are specified
- (iii) $\rho(x)$ is convex (concave)

then

$$\int_0^\infty \rho(x) \bar{G}_{T_1}(x) dx \underset{(\sum)}{\leq} \int_0^\infty \rho(x) \bar{F}(x) dx$$

$$\underset{(\sum)}{\leq} \int_0^\infty \rho(x) \bar{G}_{T_0}(x) dx.$$

PROOF: Assume $\rho(x)$ is convex. We know [1] that $\bar{F}(x) - \bar{G}_{T_0}(x)$ has exactly two sign changes, say at x_1 and x_2 . Furthermore, the order of the sign changes is $- + -$. Define a line $\ell(x)$ so that

$$\rho(x_1) = \ell(x_1) \quad \text{and} \quad \rho(x_2) = \ell(x_2).$$

Since ρ is convex, $\rho(x) - \ell(x)$ changes sign at x_1 and x_2 in the order $+$ $-$ $+$. Therefore,

$$[\rho(x) - \ell(x)][\bar{F}(x) - \bar{G}_{T_0}(x)] \leq 0$$

for all x . Integrating on x we obtain the upper bound on

$$\int_0^\infty \rho(x) \bar{F}(x) dx.$$

The lower bound follows in a similar way using $\bar{G}_{T_1}(x)$ instead of $\bar{G}_{T_0}(x)$.

The ordering of sign changes for $\rho(x) - \ell(x)$ is of course reversed when ρ is concave. ||

Using Theorem 4.1 and letting $\rho(x) = x^r$ for $r > 0$, one can obtain improved bounds on μ_r ($r > 2$) in terms of μ_1 and μ_2 .

The general problem of bounding integrals of the form

$$\int_0^\infty \phi(x, \bar{F}(x)) dx$$

when F is IHR with μ_1 and μ_2 specified seems quite difficult. However, it is possible to treat many special cases using the properties of the extremal family considered in [1]. In order to do this we repeat some of the notation in [1]. We can assume without loss of generality that $\mu_1 = 1$. It follows that μ_2 satisfies $1 \leq \mu_2 \leq 2$. Let

$$T_0 = 1 - \sqrt{\mu_2 - 1} \quad \text{and} \quad T_1 = -a_0^{-1} \log(1 - a_0)$$

where a_0 in $[0, 1]$ and T_1 simultaneously satisfy the following equations:

$$\int_0^{T_1} e^{-a_0 x} dx = \mu_1 = 1$$

$$\int_0^{T_1} x e^{-a_0 x} dx = \frac{\mu_2}{2}.$$

Let

$$\mathcal{G}_3 = \{G_T : T \geq T_1\}$$

where

$$1 - G_T(x) = \begin{cases} 1 & x < \Delta \\ e^{-a(x-\Delta)} & \Delta \leq x \leq T \\ 0 & x \geq T \end{cases} \quad T \geq T_1$$

and a, Δ ($0 \leq \Delta \leq T_0$) are chosen to satisfy the moment conditions; i.e.,

$$\int_0^{\infty} [1 - G_T(x)] dx = \mu_1 = 1$$

$$\int_0^{\infty} x[1 - G_T(x)] dx = \frac{\mu_2}{2}.$$

Let

$$\mathcal{G}_4 = \{G_T : T_0 \leq T \leq T_1\}$$

where

$$1 - G_T(x) = \begin{cases} e^{-a_1 x} & x \leq T \\ e^{-a_1 T - a_2 (x - T)} & x \geq T \end{cases} \quad T_0 \leq T \leq T_1$$

and $a_1 \leq a_2$ are chosen to satisfy the moment conditions as

before. It is shown in [1] that for $t \geq 0$

$$\inf [1 - G_T(t)] \leq 1 - F(t) \leq \sup [1 - G_T(t)]$$

where the extremums are taken over $\mathcal{H}_3 \cup \mathcal{H}_4$. These bounds have been tabulated for selected values of μ_2 ($1 \leq \mu_2 \leq 2$).

4.1 Expected Time Between Removals. If X has distribution F , then $E[\min(X, t)] =$

$$\int_0^t \bar{F}(x) dx.$$

For an age replacement policy this denotes the expected time between removals using a policy determined by t .

THEOREM 4.2. If F is IHR, $F(0) = 0$ and $\mu_1 = 1$, μ_2 are specified, then

$$(i) \quad \int_0^t \bar{F}(x) dx \geq \inf_{G_T \in \mathcal{H}_3} \int_0^t \bar{G}_T(x) dx$$

$$(ii) \quad \int_0^t \bar{F}(x) dx \leq \begin{cases} t & t \leq T_0 = 1 - \sqrt{\mu_2 - 1} \\ \int_0^t \bar{G}_T(x) dx & T_0 \leq t \leq T_1 \\ \mu_1 = 1 & t \geq T_1 \end{cases}$$

All inequalities are sharp.

PROOF: Let $F \in \mathcal{F} - (\mathcal{H}_3 \cup \mathcal{H}_4)$ where \mathcal{F} is the family of all IHR distributions with prescribed moments $\mu_1 = 1$ and μ_2 .

Let $T > T_1$ and $s(T)$ be the crossing in (Δ, T) from above of $1 - G_T$ by $1 - F$. We use crucially the fact that $s(T)$ is continuous in T (see [1] p. 52).

Case 1. $t \leq s(T_1)$. From the definition of G_{T_1} we can see that

$$\bar{F}(x) \geq \bar{G}_{T_1}(x) \text{ for } 0 \leq x \leq s(T_1)$$

and hence

$$\int_0^t \bar{F}(x) dx \geq \int_0^t \bar{G}_{T_1}(x) dx .$$

Case 2. $t \geq s(\infty)$. $\bar{F}(x)$ crosses $\bar{G}_{T_0}(x)$ from above at $s(\infty)$ and

$$\bar{F}(x) \leq \bar{G}_{T_0}(x) \text{ for } x > s(\infty) .$$


Hence

$$\int_t^\infty \bar{F}(x) dx \leq \int_t^\infty \bar{G}_{T_0}(x) dx$$

implies

$$\int_0^t \bar{F}(x) dx \geq \int_0^t \bar{G}_{T_0}(x) dx .$$

Case 3. $s(T_1) < t < s(\infty)$. Since $s(T)$ is continuous in T , choose $T \geq T_1$ and Δ ($0 \leq \Delta \leq T_0$) such that $s(T) = t$. In this circumstance $G_T \in \mathcal{H}_3$. A drawing will help.



Let

$$\phi_t(x) = \begin{cases} 1 & x \leq t \\ 0 & x > t \end{cases}$$

and define a line $\ell(x)$ such that $\ell(x_1) = 1$ and $\ell(T) = 0$.

Since clearly

$$[\phi_t(x) - \ell(x)][\bar{F}(x) - \bar{G}_T(x)] \geq 0$$

for all x ,

$$\int_0^{\infty} [\phi_t(x) - \ell(x)][\bar{F}(x) - \bar{G}_T(x)] dx =$$

$$\int_0^t \bar{F}(x) dx - \int_0^t \bar{G}_T(x) dx \geq 0.$$

This proves (i).

To prove (ii) we first note that

$$\bar{F}(x) \leq \bar{G}_{T_0}(x) = 1 \quad \text{for} \quad 0 \leq x \leq T_0$$

and hence

$$\int_0^t \bar{F}(x) dx \leq t \quad \text{for} \quad t \leq T_0,$$

is sharp.

Also,

$$\bar{F}(x) \geq \bar{G}_{T_1}(x) \quad \text{for } x \geq T_1$$

implies


$$\int_0^t \bar{F}(x) dx \leq \int_0^t \bar{G}_{T_1}(x) dx = \mu_1 = 1. \quad t \geq T_1$$

is sharp. To complete the proof we need the function $u(T)$ defined for $T_0 \leq T \leq T_1$. Let $u(T)$ be the crossing in $[T, \infty]$ from below of \bar{G}_T by \bar{F} ; $u(T)$ always exists (see [1]). We use crucially the fact that $u(T)$ varies continuously from $u(T_0) > T_0$ to $u(T_1) = T_1$.

Case 1. $t \leq u(T_0)$. Clearly $\bar{F}(x) \leq \bar{G}_{T_0}(x)$ for $x \leq u(T_0)$ implies

$$\int_0^t \bar{F}(x) dx \leq \int_0^t \bar{G}_{T_0}(x) dx.$$

Case 2. $T_0 \leq u(T_0) < t < u(T_1) = T_1$. By continuity of $u(T)$ we can choose T such that $u(T) = t$. In this circumstance $G_T \in \mathcal{U}_4$. The following drawing will help.



If $\bar{F}(x) \leq \bar{G}_T(x)$ for $x \leq t$, the proof is obvious. Hence suppose $\bar{F}(x)$ crosses $\bar{G}_T(x)$ from above at x_1 . If $x_2 = \infty$,

the proof is also obvious. Hence suppose $x_2 < \infty$. Let

$$\phi_t(x) = \begin{cases} 1 & x \leq t \\ 0 & x > t \end{cases}$$

and construct a line $\ell(x)$ so that $\ell(x_1) = 1$ and $\ell(x_2) = 0$.

Since

$$[\phi_t(x) - \ell(x)][\bar{F}(x) - \bar{G}_T(x)] \leq 0$$

for all x we have

$$\begin{aligned} & \int_0^{\infty} [\phi_t(x) - \ell(x)][\bar{F}(x) - \bar{G}_T(x)] dx \\ &= \int_0^t \bar{F}(x) dx - \int_0^t \bar{G}_T(x) dx \leq 0. \parallel \end{aligned}$$

Although the above bounds must be computed numerically, they can be easily programmed because of the simple form of the extremal distributions.

If μ_1 is the only moment specified (F IHR), then

$$\int_0^t \bar{F}(x) dx \geq \mu_1 [1 - e^{-t/\mu_1}]$$

$$\int_0^t \bar{F}(x) dx \leq \begin{cases} t & t < \mu_1 \\ \mu_1 & t \geq \mu_1 \end{cases}$$

All inequalities are sharp. The lower bound follows from the fact that $\bar{F}(x)$ crosses e^{-x/μ_1} at most once and from above. The inequality is clear for small t . If it were ever violated, then a fortiori it would also be violated for $t = +\infty$ which is impossible.

4.2 Mean Residual Life

The mean residual life of an item aged t is

$$\int_t^{\infty} \bar{F}(x) dx / \bar{F}(t).$$

If F is IHR, this ratio is necessarily nonincreasing in t .

THEOREM 4.3: If F is IHR, $F(0) = 0$ and $\mu_1 = 1$, μ_2 are specified then

$$(i) \quad \frac{\int_t^{\infty} \bar{F}(x) dx}{\bar{F}(t)} \geq \begin{cases} \mu_1 - t & t \leq T_0 \\ \inf_{G_T \in \mathcal{G}_4} \frac{\int_t^{\infty} \bar{G}_T(x) dx}{\bar{G}_T(t)} & T_0 \leq t \leq T_1 \\ 0 & t > T_1 \end{cases}$$

$$(ii) \quad \frac{\int_t^{\infty} \bar{F}(x) dx}{\bar{F}(t)} \leq \begin{cases} \mu_1 = 1 & t = 0 \\ \sup_{G_T \in \mathcal{G}_3} \frac{\int_t^{\infty} \bar{G}_T(x) dx}{\bar{G}_T(t)} & 0 < t < \infty \\ \sqrt{\mu_2 - 1} & t = +\infty \end{cases}$$

All inequalities are sharp.

PROOF: (i) Case 1. $t \leq T_0$. Since $\bar{F}(x) \leq \bar{G}_{T_0}(x)$ for $0 \leq x \leq T_0$ we see that

$$\int_t^{\infty} \bar{F}(x) dx \geq \int_t^{\infty} \bar{G}_{T_0}(x) dx = \mu_1 - t$$

is sharp and

$$\bar{F}(t) \leq \bar{G}_{T_0}(t) = 1$$

implies

$$\frac{\int_t^\infty \bar{F}(x) dx}{\bar{F}(t)} \geq \mu_1 - t.$$

Case 2. $T_0 \leq t \leq T_1$. Define $u(T)$ as in Theorem 4.2. If $t \leq u(T_0)$

$$\frac{\int_t^\infty \bar{F}(x) dx}{\bar{F}(t)} \geq \frac{\int_t^\infty \bar{G}_{T_0}(x) dx}{\bar{G}_{T_0}(t)}.$$

If $u(T_0) < t \leq u(T_1) = T_1$, then choose T and $G_T \in \mathcal{V}_4$ so that $G_T(t) = F(t)$. This can be done since $u(T)$ is continuous in T . By (ii) of Theorem 4.2 we have

$$\int_0^t \bar{F}(x) dx \leq \int_0^t \bar{G}_T(x) dx.$$

Hence

$$\frac{\int_t^\infty \bar{F}(x) dx}{\bar{F}(t)} \geq \frac{\int_t^\infty \bar{G}_T(x) dx}{\bar{G}_T(t)}$$

as desired.

Case 3. $t > T_1$. We can always choose $T = t$ and $G_T \in \mathcal{V}_3$ so that

$$\bar{G}_T(x) = 0 \text{ for } x > t \text{ and } \bar{G}_T(t) \neq 0.$$


Hence

$$\frac{\int_t^\infty \bar{F}(x) dx}{\bar{F}(t)} \geq 0$$

for $t > T_1$ is sharp.

(ii) Again we take cases.

Case 1. $t < s(T_1)$. The following drawing illustrates the situation.



Clearly

$$\int_t^\infty \bar{F}(x) dx \leq \int_t^\infty \bar{G}_{T_1}(x) dx$$

in this case and

$$\bar{F}(t) \geq \bar{G}_{T_1}(t)$$

implies

$$\frac{\int_t^\infty \bar{F}(x) dx}{\bar{F}(t)} \leq \frac{\int_t^\infty \bar{G}_{T_1}(x) dx}{\bar{G}_{T_1}(t)}.$$


Case 2. $s(T_1) < t \leq s(\infty)$. By continuity of $s(T)$, there exists $T \geq T_1$ and Δ such that $s(T) = t$ and $\bar{F}(t) = \bar{G}_T(t)$ ($G_T \in \mathcal{J}_3$).
By case 3 (i) of Theorem 4.2

$$\int_0^t \bar{F}(x) dx \geq \int_0^t \bar{G}_T(x) dx$$

implies

$$\frac{\int_t^\infty \bar{F}(x) dx}{\bar{F}(t)} \leq \frac{\int_t^\infty \bar{G}_T(x) dx}{\bar{G}_T(t)}$$

Case 3. $t \geq s(\infty)$. If $t = s(\infty)$, the following situation holds.



Hence

$$\frac{\int_t^\infty \bar{F}(x) dx}{\bar{F}(t)} \leq \frac{\int_t^\infty \bar{G}_{T_0}(x) dx}{\bar{G}_{T_0}(t)} = \sqrt{\mu_2 - 1}$$

since

$$\bar{F}(t) = \bar{G}_{T_0}(t) \text{ at } t = s(\infty).$$

Since

$$\frac{\int_t^\infty \bar{F}(x) dx}{\bar{F}(t)}$$

is nonincreasing in t we have that

$$\frac{\int_t^{\infty} \bar{F}(x) dx}{\bar{F}(t)} \leq \sqrt{\mu_2 - 1}$$

for $t \geq s(\infty)$ is sharp. \parallel

It is easy to show that if only μ_1 is specified (F IHR), then the following inequalities are sharp.

$$\frac{\int_t^{\infty} \bar{F}(x) dx}{\bar{F}(t)} \geq \begin{cases} \mu_1 - t & t \leq \mu_1 \\ 0 & t > \mu_1 \end{cases}$$

$$\frac{\int_t^{\infty} \bar{F}(x) dx}{\bar{F}(t)} \leq \mu_1.$$

4.3 Expected Time Between Failure Removals

If an item is replaced either at age t or at failure, whichever occurs first, then the expected time between failure removals will be

$$\frac{\int_0^t \bar{F}(x) dx}{F(t)}.$$

This ratio is nonincreasing in t if F is IHR. This ratio also occurs in a bound on the renewal function. Suppose F is IHR and $N(t)$ is the number of renewals in $[0, t]$ for an associated renewal process. Then it can be shown ([5], p. 8) that

$$E[N(t)] \leq t F(t) / \int_0^t \bar{F}(x) dx \leq \frac{t}{\mu_1}.$$

It is therefore of some interest to obtain bounds on this ratio.

THEOREM 4.4: If F is IHR, $F(0) = 0$ and $\mu_1 = 1$, μ_2 are specified, then

$$(i) \quad \frac{\int_0^t \bar{F}(x) dx}{F(t)} \geq \begin{cases} \inf_{G_T \in \mathcal{K}_3} \frac{\int_0^t \bar{G}_T(x) dx}{G_T(t)} & t \leq \mu_1 = 1 \\ \inf_{G_T \in \mathcal{K}_3 \cup \mathcal{K}_4} \frac{\int_0^t \bar{G}_T(x) dx}{G_T(t)} & \mu_1 \leq t < T_1 \\ \mu_1 & t \geq T_1 \end{cases}$$

$$(ii) \quad \frac{\int_0^t \bar{F}(x) dx}{F(t)} \leq \begin{cases} \infty & t \leq T_0 = 1 - \sqrt{\mu_2 - 1} \\ \sup_{G_T \in \mathcal{K}_4} \frac{\int_0^t \bar{G}_T(x) dx}{G_T(t)} & T_0 \leq t \leq T_1 \\ \sup_{G_T \in \mathcal{K}_3} \frac{\mu_1}{G_T(t)} & t > T_1 \end{cases}$$

All inequalities are sharp.

PROOF (i) Case 1. If $t \leq s(T_1)$, then $\bar{F}(x) \geq G_{T_1}(x)$ for $0 \leq x \leq t$ and

$$\int_0^t \bar{F}(x) dx \geq \int_0^t \bar{G}_{T_1}(x) dx$$


implies the result.

Case 2. If $s(T_1) \leq t \leq s(\infty)$, then there exists T such that $F(t) = G_T(t)$ where $G_T \in \mathcal{L}_3$. This follows from the continuity of $s(T)$. Hence

$$\frac{\int_0^t \bar{F}(x) dx}{F(t)} \geq \frac{\int_0^t \bar{G}_T(x) dx}{G_T(t)}$$

by (i) of Theorem 4.2. Since necessarily $s(\infty) > \mu_1$, the first bound in (i) is immediate.

Case 3. If $t \geq s(\infty)$ we use the function $v(T)$. Let $v(T)$ be the crossing in (T, ∞) from above of $1 - G_T$ by $1 - F$ if such a crossing exists. Otherwise let $v(T)$ be the right-hand endpoint of the support of F . Since $v(T)$ is continuous in T , choose T such that $v(T) = t$; i.e., we have



Since $G_T(t) = F(t)$ and clearly

$$\int_t^\infty \bar{F}(x) dx \leq \int_t^\infty \bar{G}_T(x) dx$$

we have

$$\frac{\int_0^t \bar{F}(x) dx}{F(t)} \geq \frac{\int_0^t \bar{G}_T(x) dx}{G_T(t)}.$$

Case 4. $t \geq T_1$. Since

$$\frac{\int_0^t \bar{F}(x) dx}{F(t)}$$

is nonincreasing in t , we have

$$\frac{\int_0^t \bar{F}(x) dx}{F(t)} \geq \mu_1$$

for all t . For $t > T_1$ this bound is attained by G_{T_1} . ||

(ii) Case 1. If $t \leq T_0$, then $F(t) \geq G_{T_0}(t) = 0$. Hence the inequality is trivial in this case.

Case 2. $t \leq u(T_0)$. We use the continuity of $u(T)$ again. If $t \leq u(T_0)$, then $\bar{F}(t) \leq \bar{G}_{T_0}(t)$ and

$$\int_0^t \bar{F}(x) dx \leq \int_0^t \bar{G}_{T_0}(x) dx$$


implies the result.

Case 3. If $u(T_0) \leq t < u(T_1) = T_1$, then there exists T such that $G_T(t) = F(t)$ by continuity of $u(T)$. Note $G_T \in \mathcal{S}_4$. By Theorem 4.2 (ii)

$$\frac{\int_0^t \bar{F}(x) dx}{F(t)} \leq \frac{\int_0^t \bar{G}_T(x) dx}{G_T(t)}$$

Case 4. $t \geq T_1$. There exists $G_T \in \mathcal{S}_3$ corresponding to every t

such $\bar{F}(t) \leq \bar{G}_T(t)$; i.e., \bar{G}_T is the extremal distribution furnishing an upper bound on $\bar{F}(t)$ for $t \geq T_1$. The following drawing will help.



Since $\bar{F}(x) \geq \bar{G}_T(x) = 0$ for $x > T$ we have

$$\int_0^t \bar{F}(x) dx \leq \int_0^t \bar{G}_T(x) dx ,$$

and $F(t) \geq G_T(t)$ implies

$$\frac{\int_0^t \bar{F}(x) dx}{F(t)} \leq \frac{\int_0^t \bar{G}_T(x) dx}{G_T(t)} = \frac{\mu_1}{G_T(t)} \parallel$$

$\bar{G}_T(t)$ is tabled for $\mu_1 = 1$ and μ_2 ($1 \leq \mu_2 \leq 2$) as the upper bound on $\bar{F}(t)$ for the case $t \geq T_1$.

The asymptotic expected cost per unit time under an age replacement policy where an item is replaced either at failure or at time t whichever occurs first is [cf. [7] p. 68]

$$C(t) = \frac{(c_1 - c_2) F(t) + c_2}{\int_0^t \bar{F}(x) dx}$$

($c_1 \geq c_2$) . The methods of Theorem 4.4 will also provide sharp

bounds on this function.

If only μ_1 is specified, (F IHR), then the following inequalities are sharp:

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} \geq \mu_1 \quad \text{for all } t,$$

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} \leq \begin{cases} \infty & t \leq \mu_1 \\ \frac{\mu_1}{1 - e^{-wt}} & t > \mu_1 \end{cases}$$

where w depends on t and is determined by

$$\int_0^t e^{-wx} dx = \mu_1 = 1.$$

The first inequality is attained by the exponential. The second inequality is attained by the degenerate and exponentials truncated on the right.

4.4 Expected Time Between Planned Replacements

If an item is replaced either at time t or at failure, whichever occurs first, then the expected time between planned replacements will be

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)}.$$

This ratio is always nondecreasing in t .

THEOREM 4.5 If F is IHR, $F(0) = 0$ and $\mu_1 = 1$, μ_2 are specified, then

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} \geq \begin{cases} t & 0 \leq t \leq T_0 \\ \inf_{\mathcal{U}_3} \frac{\int_0^t \bar{G}_T(x) dx}{\bar{G}_T(t)} & T_0 \leq t \leq T_1 \\ \inf_{\mathcal{U}_3} \frac{\int_0^t \bar{G}_T(x) dx}{\bar{G}_T(t)} & t > T_1 \end{cases}$$

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} \leq \begin{cases} \sup_{\mathcal{U}_3} \frac{\int_0^t \bar{G}_T(x) dx}{\bar{G}_T(t)} & 0 \leq t \leq T_0 \\ \sup_{\mathcal{U}_3} \frac{\int_0^t \bar{G}_T(x) dx}{\bar{G}_T(t)} & T_0 < t \leq T_1 \\ \infty & t > T_1 \end{cases}$$

All inequalities are sharp.

PROOF: Clearly $\bar{F}(x) \geq \bar{F}(t)$ for $0 \leq x \leq t$ implies

$$\int_0^t \bar{F}(x) dx \geq \int_0^t \bar{F}(t) dx = t \bar{F}(t)$$

for all $t > 0$. For $t \leq T_0$, the lower bound is attained with \bar{G}_{T_0} .


Now we consider cases, according as $t < s(T_1)$ or $t > s(T_1)$, where $s(T_1)$ denotes the crossing point of $\bar{G}_{T_1}(x)$ from above by $\bar{F}(x)$.

Case 1. $T_0 \leq t \leq s(T_1)$. Let $G_T \in \mathcal{G}_4$ and let $a_1(T)$ denote a solution parameter for T . [see preliminaries to this section]

It can be shown that $a_1(T)$ is a continuous function of T ([1] p. 52). Let $w(T)$ denote the crossing of \bar{G}_T by \bar{F} from above in $[0, T]$ if it exists. Clearly $w(T_1) = s(T_1)$ and

$$0 \leq w(T) \leq s(T_1).$$

Continuity of $a_1(T)$ together with log concavity of $\bar{F}(x)$ insures that for t satisfying $T_0 \leq t \leq s(T_1)$ there exists T such that $w(T) = t$ where $G_T \in \mathcal{G}_4$. A drawing will help:




Clearly we will have $\bar{F}(x) \geq \bar{G}_T(x)$ for $x \leq t$ and hence

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} \geq \frac{\int_0^t \bar{G}_T(x) dx}{\bar{G}_T(t)}.$$

Case 2. $s(T_1) < t < s(\infty)$.

Choose T and $G_T \in \mathcal{G}_3$ such that $s(T) = t$; i.e., we will have



This is possible by the continuity of $s(T)$. Hence

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} \geq \frac{\int_0^t \bar{G}_T(x) dx}{\bar{G}_T(t)}$$

by Theorem 4.2. Note that $s(T_1) \leq T_1$ so that we need only consider $G_T \in \mathcal{M}_3$ for $t \geq T_1$.

Case 3. $t > s(\infty)$

In this case $\bar{F}(x) \leq \bar{G}_{T_0}(x)$ for $x \geq t$ and therefore

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} \geq \frac{\int_0^t \bar{G}_{T_0}(x) dx}{\bar{G}_{T_0}(t)} . \parallel$$

To prove the upper bound we again take cases.

Case 1. $0 \leq t \leq T_0$. We may suppose $\bar{F}(t) < 1$ ($0 \leq t \leq T_0$) since otherwise


$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} = t$$

and this value is realized by $G_{T_0} \in \mathcal{M}_3$. Hence this is considered when we take supremums over \mathcal{M}_3 . We use crucially the continuity

of $r(T)$ where $r(T)$ is the point in (Δ, T) that \bar{F} crosses \bar{G}_T from below if such a crossing exists. Clearly $r(T_0) \geq T_0$; i.e.,




Consider now $0 \leq t \leq r(T_0)$. By continuity of $r(T)$ (see [1]) we can choose T so that $r(T) = t$. Then we have



By definition of $r(T)$, $\bar{F}(x) \leq \bar{G}_T(x)$ for $0 \leq x \leq t$ and $F(t) = G_T(t)$ where $G_T \in \mathcal{G}_3$. Hence

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} \leq \frac{\int_0^t \bar{G}_T(x) dx}{\bar{G}_T(t)}$$

Case 2. $T_0 \leq t \leq T_1$. By case 1, we may assume t satisfies $r(T_0) = u(T_0) \leq t \leq u(T_1) = T_1$. By continuity of $u(T)$ choose T and $G_T \in \mathcal{G}_4$ such that $u(T) = t$. The following drawing will help;



By Case (3) of (ii) of Theorem 4.2 we have $F(t) = G_T(t)$ and

$$\int_0^t \bar{F}(x) dx \leq \int_0^t \bar{G}_T(x) dx$$

hence

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} \leq \frac{\int_0^t \bar{G}_T(x) dx}{\bar{G}_T(t)}.$$

Case 3. $t > T_1$. In this case the upper bound is attained by $G_{T_1}(t)$; i.e.,

$$\frac{\int_0^t \bar{G}_{T_1}(x) dx}{\bar{G}_{T_1}(t)} = \mu_1/0 = +\infty. \parallel$$

If the only moment specified is the mean μ_1 (F IHR), then the following inequalities are sharp.

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} \geq \begin{cases} t & t \leq \mu_1 \\ \inf_w \frac{\int_0^t \bar{G}_w(x) dx}{\bar{G}_w(t)} & t > \mu_1 \end{cases}$$

$$\frac{\int_0^t \bar{F}(x) dx}{\bar{F}(t)} \leq \begin{cases} \sup_{\Delta} \frac{\int_0^t \bar{G}_{\Delta}(x) dx}{\bar{G}_{\Delta}(t)} & t < \mu_1 \\ +\infty & t \geq \mu_1 \end{cases}$$

where

$$\bar{G}_w(x) = \begin{cases} e^{-wx} & x < T \\ 0 & x \geq T \end{cases}$$

w and T are chosen to satisfy

$$\int_0^T e^{-wx} dx = \mu_1$$

and

$$\bar{G}_\Delta(x) = \begin{cases} 1 & 0 \leq x \leq \Delta \\ -\frac{x - \Delta}{\mu_1 - \Delta} & x \geq \Delta \end{cases}$$

($0 \leq \Delta \leq \mu_1$) . The proof is similar to previous proofs and uses the methods of [1] pp. 43-44.

Acknowledgment This paper represents an extension of part of a previous paper [1] with Albert Marshall to whom I would like to acknowledge a great indebtedness.

REFERENCES

- [1] BARLOW, R. and MARSHALL, A. (1963). Bounds for distributions with monotone hazard rate, Boeing Scientific Laboratories Document DI-82-0247.
- [2] BARLOW, R. E., MARSHALL, A. W., and PROSCHAN, F. (1963). Properties of distributions with monotone hazard rate, Ann. Math. Statist. 34 375-389.
- [3] KARLIN, S. (1959). Mathematical Methods and Theory in Games, Programming, and Economics, Addison-Wesley, Mass. 210-214.
- [4] RUSTAGI, J. S. (1957). On minimizing and maximizing a certain integral with statistical applications, Ann. Math. Statist. 28 309-328.
- [5] BARLOW, R. E. and PROSCHAN, F. (1963). Comparison of replacement policies and renewal theory implications, Boeing Scientific Research Laboratories Document DI-82-0237.
- [6] KARLIN, S. and NOVIKOFF, A. Generalized convex inequalities, Pacific Journal of Mathematics (to appear).
- [7] BARLOW, R. E. and PROSCHAN, F. (1962). Planned replacement, Studies in Applied Problems and Management Science, ed. by Arrow, Karlin, and Scarf, Stanford University Press, California.
- [8] MULHOLLAND, H. P. and ROGERS, C. A. (1958). Representation theorems for distribution functions, Proc. London Math. Soc. Vol. VIII, No. 30.
- [9] KARLIN, S. and SHAPLEY, L. S. (1953). Geometry of moment spaces, Memoir No. 12 of the Amer. Math. Soc.

BASIC DISTRIBUTION LIST
FOR UNCLASSIFIED TECHNICAL REPORTS

Head, Logistics and Mathematical
Statistics Branch
Office of Naval Research
Washington 25, D. C.

C. O., ONR Office
Navy No. 100, Box 39, F. P. O.
New York, New York

ASTIA Document Service Center
Building No. 5
Cameron Station
Alexandria, Virginia

Institute for Defense Analyses
Communications Research Division
von Neumann Hall
Princeton, New Jersey

Technical Information Officer
Naval Research Laboratory
Washington 25, D. C.

C. O., ONR Branch Office
1030 East Green Street
Pasadena 1, California
ATTN: Dr. A. R. Laufer

Bureau of Supplies and Accounts
Code OW, Department of the Navy
Washington 25, D. C.

Professor Russell Ackoff
Operations Research Group
Case Institute of Technology
Cleveland 6, Ohio

Professor Kenneth J. Arrow
Serra House, Stanford University
Stanford, California

Professor G. L. Bach
Carnegie Institute of Technology
Planning and Control of Industrial
Operations, Schenley Park
Pittsburgh, 13, Pennsylvania

Professor L. W. Cohen
Mathematics Department
University of Maryland
College Park, Maryland

Professor Donald Eckman
Director, Systems Research Center
Case Institute of Technology
Cleveland, Ohio

Professor Lawrence E. Fouraker
Graduate School of Business
Harvard University
Cambridge, Massachusetts

Professor David Gale
Department of Mathematics
Brown University
Providence 12, Rhode Island

Dr. Murray Geisler
The RAND Corporation
1700 Main Street
Santa Monica, California

Professor L. Hurwicz
School of Business Administration
University of Minnesota
Minneapolis 14, Minnesota

Dr. James R. Jackson, Director
Western Management Sciences Institute
University of California
Los Angeles 24, California

Professor Samuel Karlin
Mathematics Department
Stanford University
Stanford, California

Professor C. E. Lemke
Department of Mathematics
Rensselaer Polytechnic Institute
Troy, New York

Professor W. H. Marlow
Logistics Research Project
The George Washington University
707 - 22nd Street, N. W.
Washington 7, D. C.

BASIC DISTRIBUTION LIST
FOR UNCLASSIFIED TECHNICAL REPORTS

Professor A. Charnes
The Technological Institute
Northwestern University
Evanston, Illinois

Professor R. Radner
Department of Economics
University of California
Berkeley 4, California

Professor Murray Rosenblatt
Department of Mathematics
Brown University
Providence 12, Rhode Island

Mr. J. R. Simpson
Bureau of Supplies and Accounts
Navy Department (Code W31)
Washington 25, D. C.

Professor A. W. Tucker
Department of Mathematics
Princeton University
Princeton, New Jersey

Professor J. Wolfowitz
Department of Mathematics
Lincoln Hall, Cornell University
Ithaca 1, New York

C. O., ONR Branch Office
346 Broadway
New York 13, New York
ATTN: J. Laderman

Professor Oskar Morgenstern
Economics Research Project
Princeton University
92 A Nassau Street
Princeton, New Jersey